



## King's Research Portal

DOI:

[10.1001/jamapsychiatry.2018.2486](https://doi.org/10.1001/jamapsychiatry.2018.2486)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Fusar-Poli, P., Davies, C., & Bonoldi, I. (2018). A Case of a College Student Presenting With Mild Mental Health Problems. *JAMA Psychiatry*, 75(12), 1298-1299. <https://doi.org/10.1001/jamapsychiatry.2018.2486>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# FORECASTING RISK TO PREVENT MENTAL DISORDERS

Paolo Fusar-Poli, MD, PhD<sup>1,2,3\*</sup>, Cathy Davies, MSc<sup>1</sup>, Ilaria Bonoldi, MD, PhD<sup>4</sup>

## Affiliations

1. Early Psychosis: Interventions and Clinical Detection (EPIC) lab, Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom;
2. OASIS service, South London and Maudsley NHS Foundation Trust, London, United Kingdom;
3. Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy;
4. Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, United Kingdom.

Abstract: na

Case presentation: 250

Discussion: 613

Tables: 0

Figures: 1

Endnote reference file: Prognosis.enl

**\*Correspondence to:** Dr Paolo Fusar-Poli MD PhD, Department of Psychosis Studies, 5<sup>th</sup> Floor, Institute of Psychiatry, Psychology & Neuroscience, PO63, 16 De Crespigny Park, SE5 8AF London, UK. E-mail: [paolo.fusar-poli@kcl.ac.uk](mailto:paolo.fusar-poli@kcl.ac.uk)

**Conflict of interest Disclosures:** Dr Paolo Fusar-Poli has served on advisory boards or consulted with Lundbeck, outside of the present manuscript. No other disclosures were reported.

**Additional Contributions:** We thank the patient for granting permission to publish this information.

### **Case presentation**

A 17-year old boy was referred from the general practitioner to the local psychosis early detection clinic due to a drop in functioning and social withdrawal over the previous six months. Concerned about his increasing social isolation, his mother encouraged him to seek help. He began college six months prior but had found the workload difficult, failing his exams. During this period, he reported that he could not relate to people at college or to friends. He had no family history of mental disorders, denied any current or past use of drugs and reported no significant medical history. He was well kempt, quiet during his interview, providing short answers. He reported that he no longer enjoyed his former interests, but there were no clear signs of depressive disorders. No formal thought disorders were elicited. He believed that random people looked and talked about him when he was out in public. He was 80% convinced that this was occurring but was able to question it. He stated that these people were probably commenting on the way he looked but he did not believe these individuals meant him harm. He never acted on these thoughts. He also reported a vague feeling of perplexity and derealisation. These experiences began when he started college and continued to occur every day for up to an hour at a time (if he was outside), causing significant distress. At the Structured Clinical Interview for DSM he did not meet diagnostic criteria for any mental disorder.

### **Prognosis**

Clinical High Risk for Psychosis (CHR-P<sup>1</sup>), Attenuated Psychotic Symptoms subgroup, determined using the Comprehensive Assessment of At Risk Mental States (CAARMS)<sup>2</sup>. Increased risk of developing psychosis: 26% at 3 years (95%CI 23% - 30%)<sup>3</sup>.

What Would You Do Next?	
A	Discharge to general practitioner with no clinical recommendations
B	Pharmacotherapy with low dose atypical antipsychotics
C	Clinical follow-up for at least 3 years and indicated primary prevention
D	Treatment with omega-3 fish oil dietary supplementation

### **What to do next**

C. Clinical follow-up for at least 3 years and indicated primary prevention.

### **Discussion**


Children, adolescents and young adults (aged 8-40) who seek help at specialised early detection clinics have an enhanced risk of developing mental disorders -such as psychosis- compared to the

local age-matched general population (15%<sup>4</sup> vs 0.43%<sup>5</sup> at 3 years, respectively, Figure 1). Fully addressing their presenting problems requires forecasting their risk -i.e. probability- of developing future outcomes. The development of semi-structured psychometric interviews (prognostic tools) to assess CHR-P criteria has allowed formulating a prognosis of being at risk or not at risk for psychosis<sup>3</sup>. The original prognostic tool in this field (the CAARMS<sup>2</sup>) was developed on the basis of accumulating knowledge of the specific symptoms that may predate the onset of psychosis (defined as binary outcome). The CAARMS is transdiagnostic, allowing the presence of several comorbid mental disorders at baseline, which are frequent in these patients<sup>6</sup>. Specific training is needed to use it and its administration usually requires 2 hours in the context of a clinical assessment<sup>7</sup>. Extensive international validation has confirmed that, in people referred to psychosis early detection services, the CAARMS has an adequate prognostic performance, comparable to other prognostic instruments used in medicine<sup>3</sup>. Viceversa, it does not work well outside these samples<sup>3</sup>. Its potentials include a good ability to rule out a state of risk for psychosis, while its limitations include only a moderate ability to rule in a state of risk for psychosis (Figure 1)<sup>4</sup>. This prognostic tool has allowed one of the first preventive approaches to psychotic disorders to be implemented in psychiatry<sup>8</sup>. Several other variants of the CAARMS have been developed, with almost comparable prognostic performance<sup>3</sup>. The use of these prognostic tools in individuals seeking help at psychosis early detection clinics can impact their clinical management by: (i) informing patients about their risk of developing psychosis, (ii) establishing whether clinical monitoring for the outcome is required, and (iii) deciding whether to initiate preventive treatments or not.

Our patient was meeting the CHR-P criteria and therefore was predicted to have an enhanced risk - 26% at 3 years<sup>3</sup> (Figure 1)- of developing emerging psychotic disorders over time, compared to help-seeking individuals not meeting CHR-P criteria (who have only 1.56% risk of developing psychosis at 3 years<sup>3</sup>, Figure 1). First, we have shared with our patient the result of the prognostic test, in the context of psychoeducational support offered by psychosis early detection clinics. Informing patients about their risks is an essential component of preventive approaches in all branches of medicine. For example, CHR-P individuals accumulate several risk factors for psychosis, some of which may be potentially modifiable<sup>9</sup>. The second clinical impact of our prognostic assessment was to recommend close clinical monitoring over the ensuing 3 years, because this is the peak of risk<sup>3</sup>. Monitoring for adverse clinical outcomes is per se clinically helpful because we would be ready to intervene in the case of clinical deterioration. Finally, our patient was offered specific preventive interventions (termed indicated primary prevention) that were based on psychological therapies (cognitive behavioural therapy). These treatments aim at improving the presenting symptoms and disability<sup>10</sup> and to stop the progression to psychosis. Although psychological therapies are currently recommended (while antipsychotics are not recommended)<sup>8</sup>, their effectiveness is not fully established and other experimental therapeutics have been investigated. Over the past years, omega-3 interventions have

received high levels of interest because of their benign profile, but they were recently found not to be effective. The first ever pharmaceutical-funded randomised controlled trial in these patients (testing a phosphodiesterase-9 inhibitor) has just started and involves risk stratification, confirming that forecasting risk through prognosis is becoming a cornerstone of preventive psychiatry. Recent developments in prognostic tools might additionally allow prediction (and potential prevention) of the onset of mental disorders other than psychosis<sup>11</sup>.

**Figure 1.** Prognostic tools for forecasting emerging mental disorders (psychosis). Individuals seeking help at specialised psychosis early detection clinics have a higher risk of developing psychosis (15% at 3 years<sup>4</sup>) than the general population (0.43% at 3 years)<sup>5</sup>. Those who will meet the Clinical High Risk for Psychosis (CHR-P) criteria at the prognostic interview (CAARMS) will have only a modest increase in their level of risk for psychosis (1.8-fold, from 15% to 26%). Those not meeting the CHR-P criteria<sup>3</sup> will have a substantial decrease of their risk (9-fold, from 15% to 1.56%).



## References

1. Fusar-Poli P. The Clinical High-Risk State for Psychosis (CHR-P), Version II. *Schizophr Bull.* 2017;Jan;43(1):44-47.
2. Yung A, Yuen H, McGorry P, et al. Mapping the onset of psychosis: the Comprehensive Assessment of At-Risk Mental States. *ANZJP.* 2005;39(11-12):964-971.
3. Fusar-Poli P, Cappucciati M, Rutigliano G, et al. At risk or not at risk? A meta-analysis of the prognostic accuracy of psychometric interviews for psychosis prediction. *World Psychiatry.* 2015;14(3):322-332.
4. Fusar-Poli P, Schultze-Lutter F, Cappucciati M, et al. The Dark Side of the Moon: Meta-analytical Impact of Recruitment Strategies on Risk Enrichment in the Clinical High Risk State for Psychosis. *Schizophr Bull.* 2016;42(3):732-743.
5. Fusar-Poli P, Rutigliano G, Stahl D, et al. Deconstructing pretest risk enrichment to optimize prediction of psychosis in individuals at clinical high risk. *Jama Psychiatry.* 2016;73(2):1260-1267.
6. Fusar-Poli P, Nelson B, Valmaggia L, Yung AR, McGuire PK. Comorbid depressive and anxiety disorders in 509 individuals with an at-risk mental state: impact on psychopathology and transition to psychosis. *Schizophr Bull.* 2014;40(1):120-131.
7. Fusar-Poli P, Raballo A, Parnas J. What Is an Attenuated Psychotic Symptom? On the Importance of the Context. *Schizophr Bull.* 2017;43(4):687-692.
8. NICE. Psychosis and schizophrenia in adults: prevention and management. . 2014. <https://www.nice.org.uk/guidance/cg178/chapter/1-Recommendations - preventing-psychosis-2>. Accessed April 8th 2016.
9. Fusar-Poli P, Tantardini M, De Simone S, et al. Deconstructing vulnerability for psychosis: Meta-analysis of environmental risk factors for psychosis in subjects at ultra high-risk. *European Psychiatry.* 2017;40:65-75.
10. Fusar-Poli P, Rocchetti M, Sardella A, et al. Disorder, not just state of risk: meta-analysis of functioning and quality of life in people at high risk of psychosis. *Br J Psychiatry.* 2015;207(3):198-206.
11. Fusar-Poli P, De Micheli A, Rocchetti M, et al. Semistructured Interview for Bipolar At Risk States (SIBARS). *Psychiatry Res.* 2018;264:302-309.